



Exercises and Dry Needling for Subacromial Pain Syndrome: A Randomized Parallel-Group Trial

José L. Arias-Buría,^{*,†,‡} César Fernández-de-las-Peñas,^{†,§} María Palacios-Ceña,^{*,†}
Shane L. Koppenhaver,[¶] and Jaime Salom-Moreno^{*,†}

^{*}Department of Physical Therapy, Universidad Francisco de Vitoria, Madrid, Spain.

[†]Cátedra de Investigación y Docencia en Fisioterapia, Terapia Manual y Punción Seca, Universidad Rey Juan Carlos, Alcorcón, Spain.

[‡]Department of Physical Therapy, Hospital Universitario Gregorio Marañón, Madrid, Spain.

[§]Department of Physical Therapy, Occupational Therapy, Rehabilitation and Physical Medicine, Universidad Rey Juan Carlos, Alcorcón, Spain.

[¶]US Army-Baylor University Doctoral Program in Physical Therapy, San Antonio, Texas.

Abstract: This randomized clinical trial investigated the effectiveness of exercise versus exercise plus trigger point (TrP) dry needling (TrP-DN) in subacromial pain syndrome. A randomized parallel-group trial, with 1-year follow-up was conducted. Fifty subjects with subacromial pain syndrome were randomly allocated to receive exercise alone or exercise plus TrP-DN. Participants in both groups were asked to perform an exercise program of the rotator cuff muscles twice daily for 5 weeks. Further, patients allocated to the exercise plus TrP-DN group also received dry needling to active TrPs in the muscles reproducing shoulder symptoms during the second and fourth sessions. The primary outcome was pain-related disability assessed using the Disabilities of the Arm, Shoulder, and Hand questionnaire. Secondary outcomes included mean current pain and the worst pain experienced in the shoulder during the previous week. They were assessed at baseline, 1 week, and 3, 6, and 12 months after the end of treatment. Analysis was according to intention to treat with mixed analysis of covariance adjusted for baseline outcomes. At 12 months, 47 patients (94%) completed follow-up. Statistically larger improvements (all, $P < .01$) in shoulder disability was found for the exercise plus TrP-DN group at all follow-up periods (post: $\Delta -20.6$ [95% confidence interval (CI) -23.8 to -17.4]; 3 months: $\Delta -23.2$ [95% CI -28.3 to -18.1]; 6 months: $\Delta -23.6$ [95% CI -28.9 to -18.3]; 12 months: $\Delta -13.9$ [95% CI -17.5 to -10.3]). Both groups exhibited similar improvements in shoulder pain outcomes at all follow-up periods. The inclusion of TrP-DN with an exercise program was effective for improving disability in subacromial pain syndrome. No greater improvements in shoulder pain were observed.

Perspective: This study found that the inclusion of 2 sessions of TrP-DN into an exercise program was effective for improving shoulder pain-related disability at short-, medium-, and long-term; however, no greater improvement in shoulder pain was observed.

© 2016 by the American Pain Society

Key words: Subacromial pain syndrome, exercise, trigger point, dry needling.

Shoulder pain is a significant health problem presenting a prevalence of 25% in the general population.²⁵ Tekavec et al reported that the most prevalent diagnosis is subacromial pain syndrome.³²

The societal burden of shoulder pain is substantial with annual costs per patient estimated at €4,139 in primary health care³³ and direct costs for the treatment of shoulder disorders in the United States over \$7 billion.²⁸

Conservative treatment is the first therapeutic option for individuals with shoulder pain¹³; however, the most appropriate treatment strategy is unclear. Therapeutic exercise probably exhibits the highest level of evidence for the treatment of shoulder pain conditions including subacromial pain syndrome,^{27,30} although further trials are required.¹² In fact, the Dutch Orthopedic Association Clinical Practice Guideline for subacromial pain syndrome recommends exercise as the first therapeutic

Received April 6, 2016; Revised July 29, 2016; Accepted August 29, 2016.
The authors have no conflicts of interest to declare.

Trial registration: <http://www.clinicaltrials.gov>, NCT02338908.

Address reprint requests to César Fernández-de-las-Peñas, PT, PhD, DMSc, Facultad de Ciencias de la Salud, Universidad Rey Juan Carlos, Avenida de Atenas s/n, Alcorcón, Madrid 28922, Spain. E-mail: cesar.fernandez@urjc.es

1526-5900/\$36.00

© 2016 by the American Pain Society

<http://dx.doi.org/10.1016/j.jpain.2016.08.013>

option, but also that inactivation of trigger points (TrPs) in the shoulder be considered.⁸ TrPs are defined as hypersensitive tender spots within taut bands of skeletal muscles that are painful, elicit a referred pain, and generate motor dysfunctions.³¹ Previous studies have shown that active TrPs in the shoulder muscles reproduce symptoms suffered by subjects with subacromial pain syndrome.^{4,16}

Several therapeutic approaches, pharmacological and nonpharmacological, are proposed for the management of active TrPs, with manual therapies, TrP injections, and dry needling (DN) being among the most commonly used.⁷ Some evidence suggests that manual therapy targeting active TrPs in the shoulder musculature is effective for reducing pain and improving function in individuals with shoulder pain in the short-term,³ but there is no evidence on mid- and long-term effects. TrP-DN is defined as a "skilled intervention using a thin filiform needle to penetrate the skin that stimulates TrPs, muscles, and connective tissue for the management of musculoskeletal disorders."² Recent meta-analyses suggest that TrP-DN may be effective for neck and shoulder pain immediately after and at medium-terms.^{21,23} However, no study has investigated long-term effects of TrP-DN in patients with shoulder pain. Our objective was to conduct a randomized clinical trial to compare the 1-year effectiveness on pain and disability of the inclusion of TrP-DN into an exercise program for people with subacromial pain syndrome.

Methods

Study Design

This randomized, parallel-group clinical trial compared 2 treatments for subacromial pain syndrome: exercise only and TrP-DN plus exercise. The primary end point was 1-year improvement shoulder pain-related disability. Secondary outcomes included the current mean of shoulder pain and the worst level of pain experienced in the preceding week in the shoulder. The current report follows the Consolidated Standards of Reporting Trials extension for clinical trials.³⁵ The study was approved by the institutional review board of Universidad Rey Juan Carlos (URJC 31/2014) and the clinical trial was registered (ClinicalTrials.gov: NCT02338908).

Participants

Consecutive subjects with a diagnosis of subacromial pain syndrome from a local regional hospital (Madrid, Spain) were screened for eligibility criteria. Participants were invited to participate in the study during routine medical visits. To be eligible, they had to fulfill the following criteria: 1) unilateral nontraumatic shoulder pain, 2) shoulder pain for at least 3 months, and 3) pain intensity of at least 4 points on an 11-point numeric pain rating scale (NPRS). In our study, subacromial pain syndrome was diagnosed following the Dutch Orthopedic Association Clinical Practice Guideline in which a cluster of tests has been proposed. Therefore, patients were diagnosed when they exhibited a positive painful arc test during shoulder abduction (+ likelihood ratio

Exercises and Dry Needling for Subacromial Pain Syndrome [+LR] = 3.7; 95% confidence interval [CI], 1.9–7.0),¹⁴ and at least 2 positive of the following clinical tests: Hawkins-Kennedy test (+LR = 1.70; 95% CI, 1.29–2.26), Neer sign (+LR = 1.86; 95% CI, 1.49–2.31), empty can test (specificity = .62), drop arm test (specificity = .92), or lift-off test (specificity = .97).¹ Patients were excluded if they exhibited: 1) bilateral shoulder symptoms, 2) younger than 18 or older than 65 years, 3) history of shoulder fractures or dislocation, 4) diagnosis of cervical radiculopathy, 5) previous interventions with steroid injections in the shoulder area, 6) fibromyalgia syndrome, 7) previous history of shoulder or neck surgery, or 8) any type of intervention for the neck-shoulder area during the previous year. Additionally, because fear of needles is present in approximately 20 to 25% of subjects attending general medical practice,³⁴ we also excluded patients with fear of needles and coagulation disorders to avoid any potential risk on the experimental group. All participants signed an informed consent before their inclusion in the study.

Randomization and Masking

Patients were randomly assigned to receive TrP-DN plus exercise or exercise alone. Concealed allocation was done using a computer-generated randomized table of numbers created by a statistician who did not participate in the main trial. Individual and sequentially numbered index cards with the random assignment were prepared, folded, and placed in sealed opaque envelopes. A second external researcher opened the envelope and proceeded with allocation. Examiners blinded to group allocation obtained all outcome measures.

Interventions

Both groups received the same exercise program. No consensus exists on what exercises should be applied for individuals experiencing subacromial pain syndrome; however, it is recommended that they should be specific and of low intensity and high frequency.^{5,8} Therefore, each exercise was performed in 3 sets of 12 repetitions. Each repetition included the concentric phase after the eccentric phase of the exercise, which was slowly conducted. The program consisted of 3 exercises focusing on supraspinatus, infraspinatus, and scapular stabilizer musculature. The exercise program was taught by an experienced physical therapist in the first session and monitored in the subsequent 4 sessions, once per week during the treatment period. Each session lasted approximately 20 to 25 minutes. Participants were asked to perform the exercise program on an individual basis twice every day for 5 weeks. They were monitored during the entire treatment period for proper adherence to the exercise protocol for obtaining a 90 to 95% rate of daily practice. During the follow-up period, participants were asked to perform exercise on demand, which was monitored on subsequent follow-up assessments.

Patients allocated to the TrP-DN group also received TrP-DN to active TrPs in shoulder muscles that referred pain or reproduced shoulder symptoms during the second and

fourth treatment sessions. Therefore, patients allocated to this group received the same instructions for the exercise program in the first session, and TrP-DN during the second and fourth sessions in which participants also performed the exercise program monitored by the clinician. The muscles included in physical examination included the anterior and middle deltoid, supraspinatus, infraspinatus, teres minor and major, and subscapularis.^{4,16} Because some muscles can exhibit multiple TrPs¹¹ a clinically pragmatic approach was applied. Therefore, if multiple active TrPs were found, the clinician selected the most painful for receiving TrP-DN. Participants received TrP-DN with disposable stainless steel needles of .32 mm × 40 mm (Novasan, Madrid, Spain) that were inserted into the skin over the TrP. In this study, the fast-in and fast-out technique described by Hong¹⁷ was applied. When the active TrP was located, the overlying skin was cleaned with alcohol. The needle was inserted penetrating the skin into the TrP area until the first local twitch response was obtained. The depth of the needle depended on the muscle and ranged from 10 to 15 mm for the infraspinatus (Fig 1) or deltoid (Fig 2) muscles to 30 to 35 mm for the supraspinatus and teres major and minor muscles. Hong¹⁷ suggested that local twitch responses should be elicited during TrP-DN for a proper and successful technique. When the first local twitch response was obtained, the needling was hence moved up and down (3–5 mm vertical motions with no rotations) at approximately 1 Hz until no more local twitch responses were elicited. TrP-DN intervention had a mean duration of 5 to 10 minutes in all participants. TrP-DN was applied by a physical therapist with 10 years of clinical experience in this therapeutic approach.

Outcome Measures

Clinical records of all subjects included questions regarding the location, intensity, and duration of the symptoms, aggravating and relieving factors, and previous treatments. Pain and related disability outcomes were assessed at baseline (pre), 1 week after the last treatment (post), and 3, 6, and 12 months after the end of therapy. It has been reported that the intensity of



Figure 1. DN on active TrPs in the infraspinatus muscle. From: David G. Simons Academy, Switzerland, with permission.



Figure 2. DN on active TrPs in the deltoid muscle. From: David G. Simons Academy, Switzerland, with permission.

shoulder pain and related disability are highly associated in patients with subacromial shoulder pain²²; however, shoulder related disability is the strongest predictor for physical therapy interventions.⁶ Therefore, we decided shoulder related disability as the primary outcome. Related disability was assessed with the Disabilities of the Arm, Shoulder, and Hand (DASH) questionnaire.¹⁸ It consists of 30 items assessing: 1) degree of difficulty during the preceding week in performing physical activities because of problems in the upper extremity (21 items), 2) severity of each pain symptom, activity-related pain, tingling, weakness, and stiffness (5 items), and 3) the problem's effect on social activities, work, and sleep, and its psychological effect (4 items). Each item is answered on a 5-point scale ranging from 1 (no difficulty to perform, no symptom, or no effect) to 5 (unable to do, very severe symptom, or high effect). Responses are summed to form a raw score that is converted to a 0 to 100 scale where higher scores reflect greater related disability.¹⁸ The Spanish version of the DASH has shown high internal consistency (Cronbach $\alpha = .96$) and excellent test-retest reliability ($r = .96$).¹⁵ It has been recently reported that the minimal clinically important difference (MCID) for the DASH is 10.8 points.⁹

The secondary outcome was the intensity of shoulder pain. An 11-point NPRS (0 = no pain, 10 = maximum pain) was used to assess the patients' current level of shoulder pain and the worst level of pain experienced in the preceding week.²⁰ Mintken et al²⁹ reported that the MCID for the NPRS in individuals with shoulder pain was 1.1 points.

We also defined a successful outcome when patients observed a 50% improvement from baseline in DASH at 6- and 12-month follow-up periods.

Treatment Side Effects

Patients were asked to report any adverse event that they experienced either after the intervention or during any other part of the study. In the current study, an adverse event was defined as sequelae with any symptom perceived as distressing and unacceptable to the patient and required further treatment.

Table 1. Baseline Characteristics According to Treatment Assignment

CHARACTERISTIC	EXERCISE GROUP (N = 25)	TrP-DN WITH EXERCISE GROUP (N = 25)
Sex, male/female, n (%)	19 (76)/6 (24)	18 (72)/7 (28)
Age (y)	48 ± 6	49 ± 5
Mean years with pain	6.2 ± 1.9	5.8 ± 1.7
Side of the symptoms, n (%)		
Right	17 (68)	18 (72)
Left	8 (32)	7 (28)
Mean intensity of shoulder pain (NPRS; 0–10)	6.6 ± 1.5	7.2 ± 1.6
Mean worst pain experienced last week (NPRS, 0–10)	7.8 ± .7	8.1 ± .9
Mean DASH score (0–100)	62.0 ± 8.1	61.3 ± 6.5

Sample Size Determination

The sample size calculations were in the basis of detecting between-group differences of 10.8 points (MCID) on the main outcome measure,⁹ assuming a standard deviation of 10.5, a 2-tailed test, an α level of .05 and a desired power (β) of 90%. The estimated desired sample size was calculated to be at least 21 subjects per group. A drop out rate of 15% was expected, so 25 patients were included in each group.

Statistical Analysis

Statistical analysis was performed using SPSS software, version 21.0 (IBM Corp, Armonk, NY) and it was conducted according to intention to treat analysis for patients in the group to which they were allocated. Baseline demographic and clinical variables were compared between both groups using independent Student t-tests for continuous data and χ^2 tests of independence for categorical data. Our primary evaluation included mixed-model repeated measured analyses of covariance (ANCOVA) with time as the within-subjects factor, group as the between-subjects factor, and adjusted for baseline outcomes for evaluating between-group differences in all of the outcomes. Gender was also included in the main analysis as covariate. We used χ^2 tests to compare success rate at 6 and 12 months between groups. To enable comparison of effect sizes, standardized mean score differences (SMDs) were calculated by dividing the mean score differences between groups by the pooled standard deviation.

Results

Between January and March 2015, 60 consecutive individuals with shoulder pain were screened for eligibility criteria. Fifty (83%) satisfied all criteria, agreed to participate, and were randomly allocated into exercise (n = 25)




Figure 3. Flow diagram of patients throughout the course of the study.

or TrP-DN plus exercise (n = 25) group. Randomization resulted in similar baseline features for all variables (Table 1).

Within patients allocated to the exercise group, 2 were lost at 12 months of follow-up because they received corticosteroid injection in the shoulder, whereas 1 patient allocated to the exercise plus TrP-DN group was lost at the 6-month follow-up because of a whiplash injury. The reasons for ineligibility are shown in Fig 3, which provides a flow diagram of patient recruitment and retention. None of the participants in either group reported any other therapeutic intervention during the study, excluding the use of nonsteroidal anti-inflammatory drugs as needed but sporadically. In fact, most participants reported that they did not continue with the exercise program during the follow-up period, only sporadically when they had an exacerbation of pain. Five patients assigned to the exercise plus TrP-DN (25%) experienced muscle soreness after the first DN session,

which resolved spontaneously within 24 to 36 hours. No clinical adverse events were reported by the participants.

Adjusting for baseline outcomes, the mixed-model ANCOVA observed significant Group × Time interaction for DASH (F = 13.449; P < .001). Patients receiving exercise plus TrP-DN exhibited higher improvements in function at all follow-up periods (immediately after: Δ -20.6 [95% CI -23.8 to -17.4]; 3 months: Δ -23.2 [95% CI -28.3 to -18.1]; 6 months: Δ -23.6 [95% CI -28.9 to -18.3]; and 12 months: Δ -13.9 [95% CI -17.5 to -10.3]; all P < .001) than those receiving the exercise protocol alone (Fig 4). Between-group effect sizes were large at all follow-up periods (1.1 > SMD > 1.6) in favor of the exercise plus TrP-DN group. The inclusion of gender as covariate did not influence the results on shoulder disability (F = .861; P = .358).

The ANCOVA did not reveal significant Group × Time interactions for mean current (F = .307; P = .582) and the worst




Figure 4. Evolution of all the outcomes (pain on top and DASH on bottom) throughout the course of the study stratified according to randomized treatment assignment. Data are means and vertical lines are standard errors. ** Significant differences between groups (P < .001).

Table 2. Primary and Secondary Outcomes Before and After Intervention, 3, 6, and 12 Months According to Randomized Treatment Assignment

OUTCOME GROUP	PREINTERVENTION	POSTINTERVENTION	3 MONTHS	6 MONTHS	12 MONTHS
Mean intensity of shoulder pain (NPRS; 0–10)					
Exercise	6.6 ± 1.5 (6.0–7.2)	6.0 ± 2.4 (5.0–7.0)	3.4 ± 1.6 (2.4–4.5)	2.1 ± 1.9 (1.3–2.9)	1.6 ± 1.5 (.8–2.3)
TrP-DN	7.2 ± 1.6 (6.6–7.9)	5.9 ± 2.5 (4.9–6.9)	3.8 ± 1.5 (2.7–4.8)	1.9 ± 2.0 (1.2–2.8)	1.5 ± 1.4 (.9–2.2)
and exercise					
Worst level of shoulder pain experienced in preceding week (NPRS; 0–10)					
Exercise	7.8 ± .7 (7.4–8.2)	5.2 ± 2.7 (4.7–5.8)	3.3 ± 2.6 (2.6–4.0)	2.4 ± 2.5 (1.9–3.0)	2.0 ± 1.6 (1.5–2.5)
TrP-DN	8.1 ± .9 (7.7–8.4)	5.5 ± 2.7 (5.1–6.1)	2.9 ± 3.0 (2.2–3.6)	1.9 ± 3.3 (1.4–2.5)	1.6 ± 1.9 (1.1–2.1)
and exercise					
DASH score (0–100)					
Exercise	62.0 ± 8.1 (59.0–65.0)	43.8 ± 6.4 (41.5–46.1)	33.8 ± 12.0 (30.2–37.4)	26.9 ± 12.8 (23.2–30.7)	15.5 ± 11.1 (12.2–18.8)
TrP-DN	61.3 ± 6.5 (58.3–62.3)	23.2 ± 4.8 (20.9–25.4)	10.6 ± 3.8 (7.0–14.2)	3.4 ± 2.5 (1.5–5.4)	1.6 ± 1.8 (.6–2.8)
and exercise					

NOTE. Data are presented as mean ± SD (95% CI).

intensity ($F = .187$; $P = .668$) of shoulder pain: both groups had similar changes in shoulder pain at all follow-up periods (Table 2). No significant between-group differences were observed at any follow-up period ($P > .43$). Both groups exhibited moderate to large within-group effect sizes ($.7 > \text{SMD} > 1.4$) at 3-, 6-, and 12-month follow-ups (Fig 4). Again, these results were not significantly different according to gender (mean pain: $F = .409$; $P = .536$; the worst experienced pain: $F = .020$; $P = .888$).

A greater number of patients allocated to the exercise plus TrP-DN group experienced a successful outcome in the intention to treat analyses at 6- ($P < .001$) and 12- ($P = .047$) month follow-up periods (Table 3).

Discussion

This is the first study investigating the effect of adding TrP-DN to a standard exercise intervention for the treatment of subacromial pain syndrome. This randomized clinical trial found that inclusion of TrP-DN into an exercise program resulted in greater improvements on shoulder-related disability in subjects with subacromial pain syndrome at 3-, 6-, and 12-month follow-ups. No

significant differences in shoulder pain were observed, rather, both groups experienced similar improvements from baseline at all follow-up periods.

The Dutch Orthopedic Association Clinical Practice Guideline proposes the use of exercises for the management of individuals with subacromial pain syndrome.⁸ Further, recent systematic reviews also support the effectiveness of exercise in subacromial shoulder pain.^{27,30} Our study found that both groups experienced a similar decrease in mean current and the worst shoulder pain supporting the effectiveness of exercises for the management of subacromial pain syndrome. Within-group change scores and their 95% CIs surpassed the MCID of 1.1 points for shoulder pain²⁹ at 3, 6, and 12 months in both groups, supporting a clinical effect of the exercise program at a medium- and long-term follow-up. It is interesting to note that no changes in shoulder pain outcomes were observed in either group at 1 week postintervention. It is possible that that dosage of exercise, the exercise loading strategy, or the exercises included in our program can explain this finding. In fact, no consensus exists on which exercise program is the best for the treatment of subacromial pain disorders.^{5,8}

The novelty of this clinical trial was the application of TrP-DN for the management of subacromial pain syndrome. We observed that subjects receiving TrP-DN in addition to exercises exhibited clinically better outcomes in pain-related disability at all follow-up periods than those individuals who received the exercise program alone. In this case, between-group change scores and their 95% CIs surpassed the MCID of 10.8 points for shoulder pain-related disability⁹ in favor of the TrP-DN group at all follow-up periods, supporting a clinical effect of this intervention. This was supported by the fact that all patients allocated to the TrP-DN group attained a successful treatment outcome for pain-related disability (reduction of at least 50%) at 6 and 12 months.

There is evidence suggesting that TrPs are related to the presence of altered motor control patterns,²⁴

Table 3. Follow-Up Successful Outcomes (50% Improvement in DASH Score) According to Randomized Treatment Assignment

OUTCOME	6-MONTHS FOLLOW-UP		12-MONTHS FOLLOW-UP	
	EXERCISE ALONE (N = 25)	EXERCISE AND TrP-DN (N = 24)	EXERCISE ALONE (N = 23)	EXERCISE AND TrP-DN (N = 24)
Successful outcome	15 (60)	24 (100)	19 (82)	24 (100)
Nonsuccessful outcome	10 (40)	0 (0)	4 (18)	0 (0)

NOTE. Data are presented as n (%).

accelerated muscle fatigability,¹⁰ and increased motor activation¹⁹ in the affected and related musculature. Therefore, treatment of TrPs may effectively reduce these motor disturbances, improve motor function, and hence decrease pain-related disability. In fact, Bron et al⁴ reported that the number of active TrPs was moderately correlated with the DASH score in patients with shoulder pain, which could explain the current results. It is plausible that TrP-DN applied on the shoulder musculature at the beginning of an exercise program can improve the motor output of the shoulder stabilizers and facilitate proper shoulder function.

The results of this study should be considered according to potential strengths and limitations. Major strengths included that the study was prospectively registered, adhered to strict Consolidated Standards of Reporting Trials guidelines, used blinded outcome assessment, concealed allocation, and intention to treat analysis. Further, the trial had high retention rates at the 12-month follow-up. Among the limitations, first was that we recruited from a single clinic which may decrease the generalization of our results. Multicenter studies controlling for site and clinician effects (cluster effects) in future trials might enhance the generalizability. Second, because we did not include a no-intervention control group, we cannot be sure that the observed improvements are due to natural history of the condition, although this is unlikely because of the chronicity of the symptoms. Third, we did not include a sham needling technique, so we cannot be sure that the benefit of TrP-DN was not simply due to

the placebo effect. Nevertheless, a recent meta-analysis concluded that real needling therapy is significantly superior to sham needling irrespective of the subtype of control or sham procedure.²⁶ This can be also related to the fact that we did not assess potential expectations of the participants to receive any therapeutic intervention which could potentially affect the results. Fourth, subjects allocated to the TrP-DN group received 2 sessions on the basis of the authors' clinical experience because no current scientific data exist on the adequate frequency and dose of therapy. We do not know if a greater number of sessions would result in larger differences between interventions. Finally, because DN is applied to active TrPs, it is possible that subgroups of individuals with subacromial pain syndrome without active TrPs would not benefit from this intervention. However, we contend that these factors would be unlikely to change the overall conclusion of the study.

Conclusions

Our data indicate that the inclusion of TrP-DN into an exercise program resulted in larger clinical improvement in shoulder pain-related disability in individuals with subacromial pain syndrome. The inclusion of TrP-DN did not influence change in shoulder pain because both groups exhibited similar improvements at all follow-up periods. The current trial suggests that TrP-DN can be clinically used for improving effects of exercise programs in people with subacromial pain syndrome.

References

- Alqunae M, Galvin R, Fahey T: Diagnostic accuracy of clinical tests for subacromial impingement syndrome: A systematic review and meta-analysis. *Arch Phys Med Rehabil* 93:229-236, 2012
- APTA: Description of Dry Needling in Clinical Practice: An Educational Resource Paper. Alexandria, VA, APTA Public Policy, Practice, and Professional Affairs Unit, 2013
- Bron C, de Gast A, Dommerholt J, Stegenga B, Wensing M, Oostendorp RA: Treatment of myofascial trigger points in patients with chronic shoulder pain: A randomized, controlled trial. *BMC Med* 9:8, 2011
- Bron C, Dommerholt J, Stegenga B, Wensing M, Oostendorp RA: High prevalence of shoulder girdle muscles with myofascial trigger points in patients with shoulder pain. *BMC Musculoskelet Disord* 12:139, 2011
- Camargo PR, Alburquerque-Sendín F, Salvini TF: Eccentric training as a new approach for rotator cuff tendinopathy: Review and perspectives. *World J Orthop* 5:634-644, 2014
- Chester R, Shepstone L, Daniell H, Sweeting D, Lewis J, Jerosch-Herold C: Predicting response to physiotherapy in the treatment of musculoskeletal shoulder pain: A systematic review. *BMC Musculoskelet Disord* 14:203, 2013
- Desai MJ, Saini V, Saini S: Myofascial pain syndrome: A treatment review. *Pain Ther* 2:21-36, 2013
- Diercks R, Bron C, Dorrestijn O, Meskers C, Naber R, de Ruiter T, Willems J, Winters J, van der Woude HJ, Dutch Orthopaedic Association: Guideline for diagnosis and treatment of subacromial pain syndrome: A multidisciplinary review by the Dutch Orthopaedic Association. *Acta Orthop* 85:314-322, 2014
- Franchignoni F, Vercelli S, Giordano A, Sartorio F, Bravini E, Ferriero G: Minimal clinically important difference of the Disabilities of the Arm, Shoulder, and Hand outcome measure (DASH) and its shortened version (QuickDASH). *J Orthop Sports Phys Ther* 44:30-39, 2014
- Ge HY, Arendt-Nielsen L, Madeleine P: Accelerated muscle fatigability of latent myofascial trigger points in humans. *Pain Med* 13:957-964, 2012
- Ge HY, Fernández-de-las-Peñas C, Madeleine P, Arendt-Nielsen L: Topographical mapping and mechanical pain sensitivity of myofascial trigger points in the infraspinatus muscle. *Eur J Pain* 12:859-865, 2008
- Gebremariam L, Hay EM, van der Sande R, Rinkel WD, Koes BW, Huisstede BM: Subacromial impingement syndrome: Effectiveness of physiotherapy and manual therapy. *Br J Sports Med* 48:1202-1208, 2014
- Greenberg DL: Evaluation and treatment of shoulder pain. *Med Clin North Am* 98:487-504, 2014
- Hermans J, Luime JJ, Meuffels DE, Reijman M, Simel DL, Bierma-Zeinstra SM: Does this patient with shoulder pain have rotator cuff disease?: The Rational Clinical Examination systematic review. *JAMA* 310:837-847, 2013

18 The Journal of Pain

15. Hervás MT, Navarro Collado MJ, Peiró S, Rodrigo Pérez JL, López Matéu P, Martínez Tello I. Spanish version of the DASH questionnaire: Cross-cultural adaptation, reliability, validity and responsiveness [in Spanish]. *Med Clin* 127:441-447, 2006
16. Hidalgo-Lozano A, Fernández-de-las-Peñas C, Alonso-Blanco C, Ge HY, Arendt-Nielsen L, Arroyo-Morales M: Muscle trigger points and pressure pain hyperalgesia in the shoulder muscles in patients with unilateral shoulder impingement: A blinded, controlled study. *Exp Brain Res* 202:915-925, 2010
17. Hong CZ: Lidocaine injection versus dry needling to myofascial trigger point: The importance of the local twitch response. *Am J Phys Med Rehabil* 73:256-263, 1994
18. Hudak PL, Amadio PC, Bombardier C: Development of an upper extremity outcome measure: The DASH (Disabilities of the Arm, Shoulder, and Hand): The Upper Extremity Collaborative Group (UECG). *Am J Ind Med* 29:602-608, 1996
19. Ibarra JM, Ge HY, Wang C, Martínez Vizcaíno V, Graven-Nielsen T, Arendt-Nielsen L: Latent myofascial trigger points are associated with an increased antagonistic muscle activity during agonist muscle contraction. *J Pain* 12:1282-1288, 2011
20. Jensen MP, Turner JA, Romano JM, Fisher L: Comparative reliability and validity of chronic pain intensity measures. *Pain* 83:157-162, 1999
21. Kietrys DM, Palombaro KM, Azzaretto E, Hubler R, Schaller B, Schluskel JM, Tucker M: Effectiveness of dry needling for upper-quarter myofascial pain: A systematic review and meta-analysis. *J Orthop Sports Phys Ther* 43: 620-634, 2013
22. Kromer TO, Sieben JM, de Bie RA, Bastiaenen CH: Influence of fear-avoidance beliefs on disability in patients with subacromial shoulder pain in primary care: A secondary analysis. *Phys Ther* 94:1775-1784, 2014
23. Liu L, Huang QM, Liu QG, Ye G, Bo CZ, Chen MJ, Li P: Effectiveness of dry needling for myofascial trigger points associated with neck and shoulder pain: A systematic review and meta-analysis. *Arch Phys Med Rehabil* 96: 944-955, 2015
24. Lucas KR, Rich PA, Polus BI: Muscle activation patterns in the scapular positioning muscles during loaded scapular plane elevation: The effects of latent myofascial trigger points. *Clin Biomech* 25:765-770, 2010
25. Luime JJ, Koes BW, Hendriksen IJ, Verhaar JA, Miedema HS, Burdorf A: Prevalence and incidence of shoulder pain in the general population: A systematic review. *Scand J Rheumatol* 33:73-81, 2004
26. MacPherson H, Vertosick E, Lewith G, Linde K, Sherman KJ, Witt CM, Vickers AJ, Acupuncture Trialists' Collaboration: Influence of control group on effect size in trial of acupuncture for chronic pain: Secondary analysis of an individual patient data meta-analysis analysis. *PLoS One* 9:e93739, 2014
27. Marinko LN, Chacko JM, Dalton D, Chacko CC: The effectiveness of therapeutic exercise for painful shoulder conditions: A meta-analysis. *J Shoulder Elbow Surg* 20: 1351-1359, 2011
28. Meislin R, Sperling J, Stitik T: Persistent shoulder pain: Epidemiology, pathophysiology, and diagnosis. *Am J Orthop* 34:S5-S9, 2005
29. Mintken PE, Glynn P, Cleland JA: Psychometric properties of the shortened disabilities of the Arm, Shoulder, and Hand Questionnaire (QuickDASH) and Numeric Pain Rating Scale in patients with shoulder pain. *J Shoulder Elbow Surg* 18:920-926, 2009
30. Saltychev M, Aärimala V, Virolainen P, Laimi K: Conservative treatment or surgery for shoulder impingement: Systematic review and meta-analysis. *Disabil Rehabil* 37:1-8, 2015
31. Simons DG, Travell JG, Simons LS: *Myofascial Pain and Dysfunction: The Trigger Point Manual*. 2nd ed., Vol 1. Baltimore, Williams & Wilkins, 1999
32. Tekavec E, Jöud A, Rittner R, Mikoczy Z, Nordander C, Petersson IF, Englund M: Population-based consultation patterns in patients with shoulder pain diagnoses. *BMC Musculoskelet Disord* 13:23, 2012
33. Virta L, Joranger P, Brox JI, Eriksson R: Costs of shoulder pain and resource use in primary health care: A cost-of-illness study in Sweden. *BMC Musculoskelet Disord* 13:17, 2012
34. Wright S, Yelland M, Heathcote K, Ng SK, Wright G: Fear of needles: Nature and prevalence in general practice. *Aust Fam Physician* 38:172-176, 2009
35. Zwarenstein M, Treweek S, Gagnier JJ, Altman DG, Tunis S, Haynes B, Oxman AD, Moher D, CONSORT group, Pragmatic Trials in Healthcare group: Improving the reporting of pragmatic trials: An extension of the CONSORT statement. *BMJ* 337:2390, 2008